



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

lead to

$$\int_0^\pi \frac{d\theta}{(7 + \cos \theta)^2} = \frac{7\sqrt{3}}{576} \pi$$

so that

$$V = \frac{80\pi}{3} (8 - 3\sqrt{3}) \text{ cu. in.}$$

$$V \doteq 234.894,585,349,6 \text{ cu. in.}$$

Also solved by G. A. KNAPP, C. N. SCHMALL, GEORGE PAASWELL, H. N. CARLETON, O. S. ADAMS, and the PROPOSER.

408. Proposed by CLIFFORD N. MILLS, Brookings, South Dakota.

The ellipse $(x^2/81) + (y^2/16) = 1$ is revolved around the y -axis. Find the area of the surface generated.

SOLUTION BY CLYDE S. ATCHISON, Washington and Jefferson College.

From the given equation, we have

$$x = \frac{9}{4} \sqrt{16 - y^2}; \quad dx = -\frac{9y dy}{4 \sqrt{16 - y^2}}; \quad \text{and} \quad ds = \frac{1}{4} \sqrt{\frac{256 + 65y^2}{16 - y^2}} \cdot dy.$$

Then the area of the surface generated is

$$\begin{aligned} 2\pi \int_{y=-4}^{y=4} x \cdot ds &= 2\pi \int_{-4}^4 \left(\frac{9}{4} \sqrt{16 - y^2} \right) \left(\frac{1}{4} \sqrt{\frac{256 + 65y^2}{16 - y^2}} \right) dy = \frac{9\pi}{8} \int_{-4}^4 \sqrt{256 + 65y^2} \cdot dy \\ &= \frac{9\pi \sqrt{65}}{16} \left[y \cdot \sqrt{y^2 + \frac{256}{65}} + \frac{256}{65} \log \left(y + \sqrt{y^2 + \frac{256}{65}} \right) \right]_{-4}^4 \\ &= \frac{9\pi \sqrt{65}}{16} \left\{ \frac{144}{\sqrt{65}} + \frac{256}{65} \log \left(4 + \frac{36}{\sqrt{65}} \right) + \frac{144}{\sqrt{65}} - \frac{256}{65} \log \left(-4 + \frac{36}{\sqrt{65}} \right) \right\}, \\ &= 162\pi + \frac{144\pi}{\sqrt{65}} \log \frac{9 + \sqrt{65}}{9 - \sqrt{65}}. \end{aligned}$$

Also solved by A. M. HARDING, NELLIE L. INGALS, HORACE OLSON, C. C. YEN, G. W. HARTWELL, H. C. FEEMSTER, J. A. ECKSON, GEORGE PAASWELL, O. S. ADAMS, and PAUL CAPRON.

MECHANICS.

321. Proposed by E. J. MOULTON, Northwestern University.

The attraction, A , in any direction, due to a homogeneous sphere, on a particle at the center of the sphere, using the Newtonian law, is obviously zero. Find the error in the following method of computing A . Take cylindrical coordinates with origin at the center of the sphere; let the Z -axis extend in the direction of the attraction to be computed, and let r, θ be the polar coordinates used. Let δ be the density and R the radius of the sphere, and k the constant of gravitation. Then

$$A = \int_{z=-R}^{z=R} \int_{r=0}^{r=\sqrt{R^2-z^2}} \int_{\theta=0}^{\theta=2\pi} \frac{k\delta r dz d\theta dr}{[r^2 + z^2]^{\frac{3}{2}}} \quad (1)$$

$$= 2\pi k\delta \int_{z=-R}^{z=R} \left[\frac{-z}{(r^2 + z^2)^{\frac{1}{2}}} \right]_{r=0}^{r=\sqrt{R^2-z^2}} dz \quad (2)$$

$$= 2\pi k\delta \int_{-R}^R \left[\frac{-z}{R} + 1 \right] dz \quad (3)$$

$$= 4\pi k\delta R. \quad (4)$$